

二〇〇〇级

,

1998 8 2000 4

326

2 3

DSM-IV

1.

326

224

13.8% 44.6%

85.0% 53.6% $\chi^2=66.10$ $p<0.001$

326

21

13.8% 38.1%

85.0% 61.9%

$\chi^2=8.78$ $p=0.003$

≤

<

>

<

2.

3.

4.

5.

6.

7.

8.

118 47.97% 0.35 33.90 56.78 38.13 27.12 13.15 5

717

1

2

1		92.05%	70.01%	
			46.16%	26.92%
21.89%	16.18%	16.04%		
2		ADL		IADL
6.97%		35.62		1.80%
3		(LSIA)	29.00 21	87.14%
4		42.83%	9.29%	
		63.10%		
LSIA				
5		56.34%	36.96%	10.88%
6			57.75%	

1

2

3
4
5
6

:
:

80-90 11.26% 1979 1980 1991 15 94 10 25

[1]

[26]

SPSS10.0 8 EPI6.0

316 271 82 135 671 355

189

1995 96 -2000 01

WHO

8

KAP

FAO

WHO

UNU

1991

1.

2.

3.

4.

1.

11

46

< E-mail>

2.

4

100

GB

AOAC

3.

1991

1991

4.

2000

RNI

AI

Delphi5.0

1.

11

28

5

2.

3.

1991

4.

1

2

4 9 4 7kcal/g

3

100g

5.

2002

1996

37.62%

3.2

97 28.07%
2001

1997

- 1.
- 2.
- 3.
- 4.
- 5.

1. 447 1551 15 5
2.

1.
2. 80% 65%
96

背景

目的

方法

结果

1

()

439 75.2 150.3

109.3 152.3 37.6 81.6 24.4

2

198 29.8

108 35.4 97.5 109.3 26.5 24.3

3

1

1	HIV	AIDS			HIV	AIDS
94		1.69:1	35.54	60	20	96.8%
	62	66.0%		43	45.74%	1500
	82	87.2%		92	97.9%	

2 HIV AIDS

P <0.001

3 HIV AIDS
AIDS HIV

P <0.001

4 HIV AIDS
HIV AIDS

P <0.001

5 HIV AIDS HIV
AIDS

HIV AIDS

AIDS

HIV AIDS

HIV AIDS

A

1998 2001

4 P (P[8], P[4], P[6] and P[9]) P[8]
 P[4] 10 P-G 4 (P[8]G1, P[4]G2,
 P[8]G3 P[8]G4) 81.7% ,

1 10 12 11
 3

2

3 G G G1
 G2 G3 G4 G9 G 1999
 P 4 P (P[8], P[4], P[6] P[9])
 P[8] P[4] 10 P-G 4
 (P[8]G1, P[4]G2, P[8]G3 P[8]G4) 81.7% ,

4

46.2% 58.3%
 7.71 (2-33)
 4.22 (1-21) ,

1

3

2

- 1.
- 2.

1. WHO

2. DELPHI

3.

4.

5.

6.



1.

14

2.

3.

1.

4

12

2.

3.

4.

5

1.

5

1995

3

4

29.8%

60.9% 49.9%

3

2

9.4% 87.5%

3.

7

4.

